Archival Report

Low Dopamine D2 Receptor Increases Vulnerability to Obesity Via Reduced Physical Activity Not Increased Appetitive Motivation

Jeff A. Beeler, Rudolf P. Faust, Susie Turkson, Honggang Ye, and Xiaoxi Zhuang

ABSTRACT

BACKGROUND: The dopamine D2 receptor (D2R) has received much attention in obesity studies. Data indicate that D2R is reduced in obesity and that the TaqA1 D2R variant may be more prevalent among obese persons. It is often suggested that reduced D2R generates a reward deficiency and altered appetitive motivation that induces compulsive eating and contributes to obesity. Although dopamine is known to regulate physical activity, it is often neglected in these studies, leaving open the question of whether reduced D2R contributes to obesity through alterations in energy expenditure and activity.

METHODS: We generated a D2R knockdown (KD) mouse line and assessed both energy expenditure and appetitive motivation under conditions of diet-induced obesity.

RESULTS: The KD mice did not gain more weight or show increased appetitive motivation compared with wild-type mice in a standard environment; however, in an enriched environment with voluntary exercise opportunities, KD mice exhibited dramatically lower activity and became more obese than wild-type mice, obtaining no protective benefit from exercise opportunities.

CONCLUSIONS: These data suggest the primary contribution of altered D2R signaling to obesity lies in altered energy expenditure rather than the induction of compulsive overeating.

Keywords: Behavioral thrift, Dietary induced obesity, D2R, Reward deficiency, Running wheels, Voluntary exercise http://dx.doi.org/10.1016/j.biopsych.2015.07.009

Hypodopaminergic function, particularly reduced dopamine D2 receptor (D2R) signaling, has been implicated in obesity in both human (1–5) and animal studies (6–9), giving rise to the reward deficiency hypothesis that suggests individuals increase reward seeking—compulsive eating—to release dopamine to compensate for diminished dopamine activity (10–12). However, despite the prominence of the hypothesis in the field, it remains controversial and empirical support has not been consistent.

Although initial studies supported the link between decreased D2R and obesity, contradictory data have begun to emerge. For example, earlier imaging studies reported reduced D2R availability in obese subjects (2,5), but more recently, other studies have called this into question (13–17). Several studies have reported an association between obesity and the TaqA1 genetic variant of D2R (18–22), but an extensive, prospective study with thousands of subjects did not support that linkage (23). Consequently, the precise contribution of altered D2R to obesity remains uncertain, though presumably important.

We have recently developed an alternative hypothesis of dopamine function and propose that its primary role is to regulate behavioral energy expenditure to adapt energy allocation to the environmental energy economy, the behavioral thrift hypothesis of dopamine (24). For decades it has been well established that dopamine can regulate activity [for review (24)], including locomotor, exploratory, and voluntary activity, which is why dopamine activating drugs are termed psychostimulants. The hypodopaminergia associated with obesity (5,8,9,25–28), in contrast, would be expected to decrease activity, shifting energy balance toward greater energy conservation and storage, facilitating obesity (24). However, none of the studies of D2R and obesity in either humans or animals assess activity levels using any measure. While the role of dopamine in regulating energy expenditure in obesity or under conditions conducive to obesity has received little investigation, the contribution of sedentary lifestyle to obesity is increasingly gaining attention (29–37).

Most evidence in support of the D2R reward deficiency hypothesis of obesity is correlational, with limited direct experimental testing. Therefore, it is unclear whether altered D2R and reward processing are causes or consequences of obesity (38–40) or, alternatively, a process that co-occurs with obesity where both are mediated by another mechanism, such as altered insulin or leptin signaling (14,41–43). Research prior to the D2R reward deficiency hypothesis of obesity generally demonstrated that D2R antagonism did not alter feeding, though it altered willingness to work for food and physical

activity [e.g., (44,45); for review (46)]. More recently, two studies directly testing a causal link between D2R and obesity obtained contradictory results. In the first, Johnson and Kenny (7) used RNA interference to knockdown D2R in striatal neurons and observed acceleration in the development of dietary-induced obesity. However, the authors did not assess the potential contribution of reduced activity that may have been caused by the D2R knockdown or consider the possibility that alterations in physical activity might account for accelerated obesity, an explanation more consistent with earlier literature. In contrast, Kim et al. (42) used D2R knockouts and found the mice exhibited a lean phenotype, which the authors suggest may be mediated via D2R interactions with leptin, resulting in increased leptin signaling. Like Johnson and Kenny (7), Kim et al. (42) did not assess alterations in activity. Notably, Kim et al. (42) used D2R knockouts. Complete knockouts often induce more severe abnormalities than knockdowns, provoking the question of whether a knockdown, more consistent with reduced rather than ablated D2R signaling, would yield different results.

In the present study, we used a D2R knockdown (KD) mouse line generated via gene targeting of the D2R locus. While the complete D2R knockout mice exhibit dwarfism (47) and impaired glucose regulation (48), the KD mice are healthy and do not display dwarfism or glucose dysregulation (reported below). We set out to directly test the following causal effects: 1) if reduced D2R signaling contributes to dietary-induced obesity (DIO); and 2) the relative contribution of increased consumption and decreased energy expenditure.

METHODS AND MATERIALS

Subjects and Dietary-Induced Obesity Paradigm

Mice were group housed except during the home cage operant and indirect calorimetry tests, where they were singly housed. Mice were provided either standard chow or high-fat diet (Bio-Serv, Flemington, New Jersev; F3282, 60% fat calories) and were weighed one time per week. Half the mice had running wheels in their home cages throughout the experiment. For the primary dietary-induced obesity experiment homozygous KD mice on C57BL6/J background (originally on 129/SvJ background; backcrossed to C57BL6/J for >10 generations) were used and compared with wild-type (WT) C57BL6/J mice from Jackson Laboratories (Bar Harbor, Maine). Both male and female mice were used. The WT mice were 10 weeks of age and the KD mice were from 8 to 29 weeks of age (mean, 15 weeks) at the start of the experiment. However, there were no age effects in our measures. For the open field, both male and female KD homozygotes and littermate WT control mice all between the ages of 8 to 12 weeks (mean 11 weeks) were tested. For the indirect calorimetry, wheel acquisition, and refeeding experiments, we used littermate WT, heterozygotes, and homozygotes to examine gene-dose effects using a range of ages (12 to 28 weeks, means: WT 18.5 \pm 1; KD homozygotes, 19.4 \pm 1.7; KD heterozygotes 20.6 \pm 1.5; no significant difference, $F_{2,21}$ = .432, p = .65). All animal procedures were approved by the Institutional Animal Care and Use Committee at the University of Chicago.

Home Cage Concurrent Choice

Mice were singly housed in home cages equipped with operant levers and pellet dispenser and provided freely available high-fat diet. Mice could earn 20 mg sucrose pellets on a resetting progressive ratio schedule (i.e., after 30 minutes inactivity, the incrementing ratio reset to the beginning of the sequence). Running wheel activity was recorded in 1-minute bins. Consumption of free food was measured daily.

Glucose and Insulin Challenges

Mice were fasted for 6 hours before the procedure. A fasting glucose reading was taken before the challenge (time 0) and then either 1 g/kg dextrose or .75 U/kg of human insulin was administered (intraperitoneal) for the glucose and insulin challenges, respectively. Subsequent glucose blood levels were determined at 15, 60, 90, and 120 minutes. Glucose was measured using Accu-check (Roche, Basel, Switzerland) with blood from the tail vein.

Further information on methods is available in Supplement 1.

RESULTS

Validation of D2R Knockdown Mouse Line

In mice heterozygous and homozygous for the knockdown, D2R messenger RNA was reduced to 55% and 3%, respectively, of WT littermates (Figure S1 in Supplement 1 for details). The KD mice exhibited a normal range body weight (below) and exhibited no baseline impairment in glucose regulation (below). Homozygote KD mice were used throughout unless otherwise noted.

KD Mice Do Not Exhibit Greater Weight Gain in DIO but Fail to Benefit from Voluntary Exercise Opportunities

There was no significant genotype difference between WT and KD mice in initial weight. Both WT and KD mice exhibited substantial weight gain across 29 weeks on a high-fat diet (genotype main effect, $F_{1,33} = 2.614$, p = .11; Figure 1A), showing an approximately 100% increase in body weight from initial to final weights (Figure 1B). However, provision of running opportunities dramatically decreased weight gain and was protective against obesity in WT mice (wheel main effect, $F_{1.31} = 40.95$, p < .001; Figure 1A,B) but not KD mice where no difference was observed between those with and without running wheels (genotype \times wheel interaction, $F_{1,31} =$ 27.19, p < .001). In this and other experiments, we used both male and female mice. Throughout we observed few sex differences, reviewed in Supplement 1. Measures and analyses described below were all conducted with these mice unless otherwise noted.

KD Mice Show Decreased Voluntary Activity Not Increased Consumption

Following 29 weeks exposure to the high-fat diet, the grouphoused mice were individually housed during a 2-week test period in a home cage equipped for concurrent choice task with free access to the same high-fat diet and continued



Figure 1. Effect of running wheels on wild-type (WT) and dopamine D2 knockdown (D2KD) mice in dietaryinduced obesity. (A) Body weights across experiment (weights taken $1 \times$ /week). (B) Percent change from initial to final weight, sexes shown separately. (C) Total caloric consumption during a 14-day home cage concurrent choice with freely available high-fat diet and sucrose via operant responding (body weight [b.w.]). (D) Total daily wheel running across 14day period. (E) Average daily running across 14-day period. **p < .01, ***p < .001. WT: n = 16/group; D2KD: n = 8/group. N.S., not significant.

access to running wheels in the wheel groups. WT mice with access to wheels consumed significantly more food compared with WT mice without access to wheels (genotype × wheels, $F_{1,29} = 10.02$, p < .01; Figure 1C), suggesting the availability of an exercise wheel increases both expenditure and intake but maintaining a lower body weight with less storage. In contrast, the availability of a running wheel had no effect on consumption in the KD mice. Regardless of wheel status, the KD mice consumed the same amount as WT mice without wheel access. To assess whether the KD mice may have difficulty feeding in an energy-depleted state, we fasted the mice and measured refeeding. After an initial 60 minutes of refeeding, there was no difference between genotypes (Supplement 1).

KD mice provided running wheels showed dramatically lower activity compared with WT mice ($F_{1,12} = 16.3$, p < .01; Figure 1D,E). These data indicate that the KD mice expended dramatically less energy on running, suggesting that the primary vulnerability to obesity associated with a reduction in D2R signaling is decreased physical activity rather than increased appetitive motivation.

KD Mice Have Reduced Bouts, Duration, and Running Speed

Figure 2A shows average running activity in the WT (blue) and KD (red) mice in 1-minute bins across the entire experiment. Though the KD mice ran considerably less, they did run. The circadian pattern was similar between genotypes and not disturbed in the KD mice. Plotting the data across a 24-hour period averaged across the 14-day experiment (Figure 2B), both genotypes showed a similar onset of running at lights out (6:00 PM) with

scattered episodes of running during the inactive period (6:00 AM). We see greatly reduced wheel running by KD mice in both the active and inactive periods. To characterize activity patterns in greater detail, we performed a bout analysis with a bout defined as consecutive minute bins with activity (Table 1). A single 1minute bin without activity terminated a bout. The greatest difference between genotypes appeared in the total number of bouts (Figure 2C1), with the KD mice showing dramatically fewer total bouts of running ($F_{1,13} = 190.49$, p < .001), consistent with much greater interbout intervals (F_{1,13} = 90.21, p < .001; Figure 2C4). KD mice also showed significantly reduced bout duration ($F_{1,13} = 5.45$, p < .05), on average running about half as long as WT mice, though this was less pronounced. The number of turns in a minute of running reflects both the average distance and the speed of running. The KD mice ran significantly slower than WT ($F_{1.13} = 33.54$, p < .001; Figure 2C3). Overall, the KD mice showed a consistent pattern of reduced running wheel activity on all measures, having fewer bouts of running that were, on average, shorter and slower. In a separate experiment, we tested initial acquisition of running behavior. All genotypes showed similar acquisition of running behavior with a trend toward a gene-dose response where reduced D2R reduced running; however, this did not reach significance, suggesting the marked difference observed here emerges over time and exposure to the running wheel (Figure S2 in Supplement 1).

KD Mice Show Reduced Exploratory Activity in the Open Field

It is possible that the observed reduction in wheel running in KD mice was caused by impaired running rather than



Figure 2. Wheel activity patterns in wild-type (WT) and dopamine D2 knockdown (D2KD) mice. (A) Total number of wheel turns in 1-minute bins across the entire 14-day experiment. (B) Wheel turns in 1-minute bins across circadian cycles of a 24-hour period, averaged across experiment. (C) Bout analysis, showing (C1) average daily number of running bouts, (C2) average duration of bouts, (C3) average speed of running (turns/min), and (C4) the average interbout interval. $^{*}p < .05$, $^{**}p < .01$. WT: n = 11; D2KD: n = 6. N.S., not significant.

decreased energy expenditure; the KD mice performed poorly on the accelerating rotarod (data not shown). Reduction of D2R may affect multiple functions, including both motor learning and motivational regulation of both energy intakeappetitive motivation-and energy expenditure, voluntary exercise. The question is the degree to which reduced wheel running arises as a consequence of motor learning deficits versus altered regulation of energy expenditure. The bout analysis (Figure 2) suggests a primary motivation/expenditure deficit. During both active and inactive cycles, the KD mice ran much slower than WT mice; however, during the active cycle, the KD mice increased their speed and ran at speeds comparable with the WT mice during the inactive period, suggesting the KD mice were capable of running WT speeds during the inactive cycle but did not. To assess activity in a nonskilled paradigm, we tested the genotypes in the open field using naive mice not exposed to wheel running or high-fat diet. Consistent with decreased energy expenditure, the KD mice showed reduced open field activity ($F_{1,16} = 4.15$, p =.0584; genotype \times block, $F_{1,628}$ = 4.15, p < .001; Figure 3A) and significantly increased resting time ($F_{1,16} = 4.552$, p < .05; Figure 3B). Moreover, they exhibited a trend toward fewer ambulatory episodes ($F_{1,16} = 3.42$, p = .08; Figure 3C); however, there was no difference in average velocity during ambulation ($F_{1,16} = .164$, p = .69; Figure 3D). Thus, the KD mice showed decreased energy expenditure in the unskilled open field test with no evidence of motor slowing (i.e., their movement velocity was the same).

KD Mice Show Increased Metabolism and Decreased Activity in Indirect Calorimetry

To assess basal metabolism (in the absence of exercise wheels and on standard chow), we tested a naive group of male mice, including heterozygotes for the KD, using indirect calorimetry, measuring metabolic rate, consumption, and activity. All mice were approximately 24 weeks of age (no age difference between genotypes), weighing 26 to 29 grams, with a trend ($F_{2,21} = 3.069$, p = .06) toward the homozygote KD mice weighing less (homozygote mean, 26.7 \pm .86 vs. 28.9 \pm .97 for WT). Percent of body mass composed of fat, as assessed by dual-energy x-ray absorptiometry (DEXA) (Supplement 1), was not significantly different between genotypes ($F_{2,21} = 2.17$, p = .13; homozygotes, 16.05 \pm .85; heterozygotes, 18.48 \pm .90; WT, 16.45 \pm .91). Consistent with findings reported above, there was no difference in food consumption normalized to body weight between the genotypes ($F_{2,21} = .157$, p = .85; Figure 4A). Surprisingly, however, the KD mice showed increased basal metabolism ($F_{2,21}$ = 7.64, p < .01; Figure 4B). Although the mechanism by which this might arise is not clear (see Discussion), it indicates that the observed vulnerability to weight gain due to reduced voluntary activity does not arise from lower basal metabolic rate. Consistent with reduced voluntary (wheel running) and exploratory (open field) activity, the KD mice showed a significant reduction in basal activity ($F_{2,21} = 6.872$, p < .01; Figure 4C). Interestingly, the heterozygotes were similar to WT



Figure 3. Open field activity in wild-type (WT) and dopamine D2 knockdown (D2KD) mice. (A) Ambulatory distance (cm) in 5-minute blocks across a 60-minute testing session, averaged across three consecutive sessions. (B) Time resting (no ambulation) in 5-minute blocks. (C) Average number of ambulatory blocks per 5-minute block. (D) Average ambulatory velocity across entire session. *p < .05. n = 8. N.S., not significant.

in terms of basal metabolic rate (Figure 4B); however, in terms of basal activity, measured in the calorimetry chambers, they were closer to the KD homozygotes, suggesting that regulation of basal locomotor activity might be particularly sensitive to D2R signaling. There were no significant differences in vertical activity counts ($F_{2,21} = 1.73$, p = .20; Figure 4D).

KD Mice Do Not Show Enhanced Appetitive Motivation in Concurrent Choice

Mice were housed in home cage operant chambers with highfat diet freely available in the cage and sucrose pellets available through operant responding on a resetting progressive ratio, where the ratio reset after 30 minutes of inactivity on the levers [Beeler et al. (24)]. Because the mean body weight between groups differed, results (where appropriate) are normalized to body weight. No differences were observed in average daily sucrose pellets earned between groups ($F_{1,35} =$ 1.82, p = .18), though the KD mice with running wheel access exhibited a trend toward decreased sucrose consumption (genotype \times wheels, $F_{1,29} = 2.9$, p = .09; Figure 5A). The percentage of total caloric intake derived from sucrose was not significantly different between genotypes ($F_{1,29} = 2.74$, p = .11), but provision of a running wheel decreased sucrose as a percentage of intake across both genotypes ($F_{1,29} = 21.6$, p < .001; Figure 5B). There were no significant genotype differences in the overall amount of active and inactive lever pressing (active lever, $F_{1,29} = 1.59$, p = .21; Figure 5C,D). Average breakpoint was not different between genotypes $(F_{1,29} = 2.22, p = .14;$ Figure 5E); however, wheel access significantly reduced breakpoint in both genotypes ($F_{1,29}$ = 7.81, p < .01). There was no genotype effect on meal size (effectively, normalized breakpoint; $F_{1,29} = 1.51$, p = .22; Figure 5F) but a trend toward the KD mice with wheel access eating smaller meals (genotype \times wheels, $F_{1,29} = 4.12$, p =.051). Similarly, there was no genotype effect on the number of bouts of pressing for sucrose ($F_{1,29} = 3.0, p = .09$; Figure 5G), although again the KD mice with wheel access showed a trend toward reduced meal number ($F_{1,29} = 3.41$, p = .075; Figure 5G). In the home cage progressive ratio, mice can modulate the average cost of sucrose pellets by shifting between longer, more costly episodes of pressing and shorter, less costly (but more frequent) episodes (meal size, Figure 5F; number of daily meals, Figure 5G). Consistent with breakpoint, provision of a running wheel decreased cost per pellet in both genotypes ($F_{1,29} = 4.81$, p < .05; Figure 5H).

In summary, the KD mice did not exhibit increased appetitive motivation in a concurrent choice task. The provision of a running wheel decreased effort toward sucrose in both genotypes, an effect more pronounced in the KD mice. In the KD mice, this cannot be attributed to a motor impairment, as the KD animals without wheel access pressed the same amount overall as WT mice. In the WT mice with wheel access, the reduced breakpoint might be attributed to reduced body weight; however, these mice consumed more free chow, suggesting that the reduction in sucrose seeking is motivational rather than reflecting overall decreased consumption. These data suggest that wheel running can decrease effort expended toward sucrose, possibly suggesting a competition in the allocation of energy between two sources of reward. This competition may be more pronounced in the KD mice that, overall, exhibited less behavioral energy expenditure. Importantly, these data also indicate that although the KD mice ran dramatically less than WT mice, this minimal running can impact their appetitive energy expenditure.

	WT		D2KD	
	Male	Female	Male	Female
Active Phase				
Number of bouts	42.5 ± 1.18	39.8 ± .69	13.28 ± .93	9.53 ± 1.11
Duration	7.12 ± .12	9.41 ± .14	4.52 ± .17	4.64 ± .47
Turns	563.55 ± 11.6	837.49 ± 16.1	134.48 ± 7.77	142.53 ± 22.07
Speed	64.38 ± .68	66.25 ± .61	19.01 ± .55	14.98 ± .9
Inactive Phase				
Number of bouts	6.33 ± .52	6.81 ± .43	2.23 ± .32	1.42 ± .31
Duration	2.27 ± .11	9.41 ± .14	2.26 ± .27	1.5 ± .13
Turns	66.2 ± 7.4	113.9 ± 9.9	42.8 ± 11.8	11.35 ± 5.24
Speed	17.14 ± 1.05	$22.43~\pm~.95$	8.2 ± .96	4.1 ± 1.13

Table 1. Bout Analysis by Genotype, Sex, and Circadian Phase

Duration in minutes, speed in revolutions/minute.

D2KD, ; WT, wild-type.

KD Mice Glucose Regulation Unaltered by Provision of Running Wheels

It is well known that exercise can reduce risk of type 2 diabetes, including in mice on high-fat diets. We administered both glucose and insulin challenges to the mice to assess whether different patterns of voluntary activity could be observed in glucose regulation. There was no main effect of genotype on fasting glucose levels ($F_{1,34} = .71$, p = .40; Figure 6A,B). The opportunity for voluntary exercise significantly reduced fasting glucose levels (wheel main effect, $F_{1,34} = 8.49$, p < .01), an effect more pronounced in the WT mice (genotype × wheel, $F_{1,34} = 22.26$, p < .001). In both the glucose and insulin challenges, a difference was observed between WT mice with

and without wheel access (Figure 6C,D). WT mice with wheel access showed significantly improved glucose clearance (WT area under the curve [AUC], $F_{1,25} = 8.59$, p < .01; Figure 6C) and a trend toward greater sensitivity to insulin (WT AUC, $F_{1,25} = 2.66$, p = .11; Figure 6D). In contrast, no difference was observed in the KD mice as a result of having access to running wheels (KD AUC: glucose, $F_{1,13} = .04$, p = .84; insulin, $F_{1,13} = .12$, p = .72; Figure 6E,F).

DISCUSSION

Reduction of D2R in the KD mice did not increase obesity in a standard DIO paradigm. The lack of an observed increase in



Figure 4. Indirect calorimetry. (A) Average daily consumption normalized to body weight (b.w.), (B) metabolic rate (kilocalorie [kcal]/hr/kg b.w.), (C) horizontal activity counts, and (D) vertical activity counts for active and inactive periods for dopamine D2 knockdown (D2KD) homozygotes (hom) (red), heterozygotes (het) (pink), and wild-type (WT) littermates (blue). **p < .01. WT: n = 7; D2KD: n = 9, hom; n = 8, het. N.S., not significant.



Figure 5. Home cage concurrent choice with freely available high-fat diet. (A) Average daily sucrose pellets earned per gram of body weight (b. w.). (B) Percentage of total daily kilocalorie (kcal) intake derived from sucrose. (C) Average daily active lever presses (LP) for sucrose normalized to body weight. (D) Average daily inactive lever presses (not normalized). (E) Average breakpoint for bouts of sucrose seeking. (F) Average size of bouts of sucrose consumption normalized to body weight. (G) Average daily number of bouts of sucrose seeking. (H) Average cost per pellet in lever presses. *p < .05, ***p < .001. WT: n = 16/group; D2KD: n = 8/group. D2KD, dopamine D2 knockdown; N.S., not significant; WT, wildtype.

consumption is inconsistent with the reward deficiency hypothesis, suggesting that reduced D2 receptor expression does not drive compulsive appetitive behavior. This is further supported in the home cage concurrent choice task where we did not observe greater effort exerted for sucrose pellets, a preferred food, in the KD mice; in fact, both genotypes exhibited reduced effort for sucrose with the provision of wheel access, an effect more pronounced in the KD mice, despite their low levels of running.

In WT mice, provision of running wheels - an opportunity for voluntary exercise-dramatically reduced weight gain under a high-fat diet that normally induces obesity, as observed previously (34,49-55). In contrast, the KD mice obtained no protective benefit from running wheels. Though the KD mice did run in the wheels and demonstrated similar patterns of circadian activity, they ran many fewer bouts, of shorter duration and slower speed, consistent with decreased energy expenditure. Though we cannot conclusively rule out the possibility of motor impairment in KD mice that diminished their activity on the running wheel, the problem appeared to lie in motivation to run rather than ability [see also (56)]. The KD mice showed reduced wheel running speed, but the circadian patterns of activity remained similar. For example, during the inactive phase, the KD mice ran slower than WT mice (Table 1). However, during the active cycle, while the KD mice again ran much slower than WT mice, they achieved speeds approximating those observed in the WT mice during the

inactive phase, suggesting their reduced speed, at least during the inactive cycle, did not arise from inability to run. Critically, we also observed a reduction in activity in both the open field and indirect calorimetry (basal activity), both of which measure unskilled activity. Overall, our data suggest that the KD mice exhibited reduced physical activity.

Our finding of increased metabolic rate in the KD mice was unexpected but consistent with a prior observation of increased metabolic rate (also measured using indirect calorimetry) in D2R knockout mice (42). In that study, D2R knockout resulted in a lean phenotype, including reduced consumption, which the authors attributed to altered leptin signaling in the hypothalamus. Though KD mice may be expected to show a less severe phenotype than knockout mice, the increased metabolic rate we observed may be attributable to a similar mechanism. Though we cannot speculate on the relationship between increased metabolic rate and decreased physical activity in the KD mice, what is clear is that reduced D2R in both these lines does not increase appetitive motivation but rather induces complex effects on energy regulation, apparently increasing basal metabolism while reducing physical activity, though Kim et al. (42) did not report locomotor activity.

Finally, the glucose and insulin challenge data indicate that such differences in activity can affect glucose regulation. The provision of running opportunities to WT mice increased their glucose clearance and insulin sensitivity, while the reduced



Figure 6. Effect of wheels on glucose regulation in wild-type (WT) and dopamine D2 knockdown (D2KD) mice. Fasting (6 hour) blood glucose levels for (**A**) male and (**B**) female mice. (**C**, **E**) Glucose challenge (dextrose, 1 g/kg total body mass, intraperitoneal) for WT (blue) and D2KD (red). Offset: Area under the curve (AUC). (**D**, **F**) Insulin challenge (.5 U/kg total body mass, intraperitoneal) for WT (blue) and D2KD (red). Offset: Area under the curve (AUC). (**D**, **F**) Insulin challenge (.5 U/kg total body mass, intraperitoneal) for WT (blue) and D2KD (red). Offset: AUC. **p < .01. WT: n = 16/group; D2KD: n = 8/group. N.S., not significant.

voluntary activity in the KD mice precluded this potentially protective effect. Though in one sense this seems obvious, as the KD mice exhibited dramatically reduced activity, the amount of activity necessary to yield an ameliorative effect on glucose metabolism is not well established. Thus, it could have been that even minimal voluntary exercise, as exhibited by the KD mice, might have altered glucose regulation, as it apparently altered appetitive motivation in the concurrent choice task (Figure 5). However, these data suggest that activity is reduced in the KD mice sufficiently to preclude potential protective effects against metabolic disorder. Additionally, the glucose and insulin challenge data indicate that the KD mice, unlike the D2R knockouts (48), do not exhibit impaired glucose homeostasis compared with WT mice.

Together, these data suggest that the role of reduced D2 receptor in obesity lies not in increasing appetitive motivation and generating compulsive eating but rather in altering activity and energy expenditure, favoring reduced behavioral expenditure of energy. More broadly, these data are consistent with a fundamental role for dopamine in regulating behavioral energy expenditure, as long suggested by Salamone *et al.* [for review (46)]. Over decades, they have demonstrated that reduced dopamine diminishes the willingness to work for reward as well as locomotor activity without altering free-feeding food preferences (46). The thrift hypothesis builds

upon the insight of Salamone et al. (46), suggesting dopamine not only regulates behavioral energy expenditure along a conserve-expend axis but also regulates the degree to which prior reward biases behavioral choice with decreased dopamine inducing greater exploitation of prior reward information [for review (24)]. In the present study, we observed no change in appetitive behavior and choice in the home cage concurrent choice as a consequence of reduced D2R signaling alone; however, in the context of an enriched environment with access to a running wheel, we observed a decrease in pursuit of sucrose among both genotypes, more pronounced in the KD mice. We speculate that in an enriched environment with more rewarding options for energy expenditure, decreased D2R induces a regime of energy conservation in which energy expenditure has to be divided rather than increased, though testing this hypothesis remains for future studies.

Several cautions are in order. First, the D2R knockdown is global and constitutive. Thus, we cannot isolate its effects to specific neural substrates, nor can we rule out compensations. However, there are two primary views of reduced D2R function in obesity. In one, the reduction precedes and causes obesity, perhaps as a consequence of genetic variance [e.g., (4,13,20,57,58)]. This potential mechanism is analogous to our genetic knockdown, as one might expect that similar compensations might occur in individuals with genetically

reduced D2R function. The other proposed mechanism is that obesity causes a reduction in D2R, which then further contributes to and maintains obesity. The development of obesity, however, is a gradual process and presumably so is the reduction in D2R. Consequently, it is likely that even with this second proposed mechanism, compensations would be induced by reduced D2R expression.

Perhaps more importantly, a recent human imaging study by Guo et al. (16) suggested that obesity-induced changes in D2R binding potential are not uniform, demonstrating a positive correlation between D2R binding in the dorsal striatal regions and both body mass index and opportunistic eating, while the ventral striatum was negatively correlated, though this latter finding did not reach significance. These data suggest that obesity may induce region-specific alterations in D2R, which may differentially contribute to behavior. Thus, our global knockdown may obscure subtleties in obesityinduced regulation of D2R and its behavioral effects. Importantly, in the Guo et al. (16) study, the strongest finding was that lateral striatal D2R binding increased with body mass index. The present findings highlight the importance of assessing energy expenditure, almost without exception neglected, when investigating dopamine-related contributions to obesitv.

Finally, we cannot assess the exact degree to which potential motor impairments induced by D2R knockdown contributed to decreased wheel-running behavior. The expenditure of calories does not require proficient but rather persistent running, which the KD mice did not exhibit. Crucially, we observed decreased activity in measures of nonskilled voluntary activity as well, including the open field and basal activity as measured in the calorimetry experiments.

Over the last two decades, there has been much focus on diet as the root cause of the increase in obesity rates, the socalled obesogenic Western diet. However, people have also become considerably more sedentary over recent decades, which is believed to contribute to obesity and risk of metabolic and cardiovascular disease (59–66). Despite evidence for central control of physical activity level by multiple substrates [e.g., (67–69)], the neural systems regulating voluntary physical activity remain poorly understood (68). Here, we show that reduced D2R expression, frequently believed to contribute to obesity, significantly alters patterns of energy expenditure with little effect on appetitive behavior, suggesting a key role of dopamine dysregulation in obesity might lie in altered behavioral energy regulation.

The thrift hypothesis of dopamine suggests that altered dopamine function will shift voluntary energy expenditure, with the reduced dopamine function associated with obesity paradoxically favoring behavioral energy conservation, i.e., reduced physical activity. Such maladaptive regulation of behavioral energy expenditure may contribute not only to promoting obesity but also to helping explain the motivational resistance often incurred by exercise programs aimed at reducing weight in obese and overweight individuals.

ACKNOWLEDGMENTS AND DISCLOSURES

This work was supported by the National Institute on Drug Abuse, R01DA25875 (JAB); the National Institute of Diabetes and Digestive and

Kidney Diseases, R56DK088515 (XZ); and the National Institutes of Health DK20595 (HY).

We thank Jared T. Hinkle for technical assistance. We also thank Claus Kemkemer from the Long lab and Wei Zhang from the Kronfrost lab.

All authors report no biomedical financial interests or potential conflicts of interest.

ARTICLE INFORMATION

From the Department of Psychology (JAB), Queens College and the Graduate Center, City University of New York, Flushing, New York; and Department of Neurobiology (JAB, ST, XZ), Interdisciplinary Scientist Training Program (RPF), Janelia Research Campus and Department of Medicine (HY), University of Chicago, Chicago, Illinois.

Authors JAB and RPF contributed equally to this work.

Address correspondence to Jeff A. Beeler, Ph.D., Queens College and the Graduate Center, City University of New York, Department of Psychology, 65-30 Kissena Blvd, Flushing, NY, 11367; E-mail: jbeeler@qc.cuny.edu.

Received Nov 12, 2014; revised Jul 15, 2015; accepted Jul 16, 2015.

Supplementary material cited in this article is available online at http://dx.doi.org/10.1016/j.biopsych.2015.07.009.

REFERENCES

- van de Giessen E, Celik F, Schweitzer DH, van den Brink W, Booij J (2014): Dopamine D2/3 receptor availability and amphetamineinduced dopamine release in obesity. J Psychopharmacol 28: 866–873.
- de Weijer BA, van de Giessen E, van Amelsvoort TA, Boot E, Braak B, Janssen IM, et al. (2011): Lower striatal dopamine D2/3 receptor availability in obese compared with non-obese subjects. EJNMMI Res 1:37.
- Volkow ND, Wang G-J, Telang F, Fowler JS, Thanos PK, Logan J, et al. (2008): Low dopamine striatal D2 receptors are associated with prefrontal metabolism in obese subjects: Possible contributing factors. Neuroimage 42:1537–1543.
- Stice E, Spoor S, Bohon C, Small DM (2008): Relation between obesity and blunted striatal response to food is moderated by TaqlA A1 allele. Science 322:449–452.
- Wang GJ, Volkow ND, Logan J, Pappas NR, Wong CT, Zhu W, et al. (2001): Brain dopamine and obesity. Lancet 357:354–357.
- Huang X-F, Zavitsanou K, Huang X, Yu Y, Wang H, Chen F, et al. (2006): Dopamine transporter and D2 receptor binding densities in mice prone or resistant to chronic high fat diet-induced obesity. Behav Brain Res 175:415–419.
- Johnson PM, Kenny PJ (2010): Dopamine D2 receptors in addictionlike reward dysfunction and compulsive eating in obese rats. Nat Neurosci 13:635–641.
- Geiger BM, Haburcak M, Avena NM, Moyer MC, Hoebel BG, Pothos EN (2009): Deficits of mesolimbic dopamine neurotransmission in rat dietary obesity. Neuroscience 159:1193–1199.
- Geiger BM, Behr GG, Frank LE, Caldera-Siu AD, Beinfeld MC, Kokkotou EG, Pothos EN (2008): Evidence for defective mesolimbic dopamine exocytosis in obesity-prone rats. FASEB J 22:2740–2746.
- Blum K, Liu Y, Shriner R, Gold MS (2011): Reward circuitry dopaminergic activation regulates food and drug craving behavior. Curr Pharm Des 17:1158–1167.
- 11. Volkow ND, Wise RA (2005): How can drug addiction help us understand obesity? Nat Neurosci 8:555–560.
- 12. Kenny PJ (2010): Reward mechanisms in obesity: New insights and future directions. Neuron 69:664–679.
- Eisenstein SA, Antenor-Dorsey JAV, Gredysa DM, Koller JM, Bihun EC, Ranck SA, et al. (2013): A comparison of D2 receptor specific binding in obese and normal-weight individuals using PET with (N-[(11)C]methyl)benperidol. Synapse 67:748–756.
- Dunn JP, Kessler RM, Feurer ID, Volkow ND, Patterson BW, Ansari MS, et al. (2012): Relationship of dopamine type 2 receptor binding potential with fasting neuroendocrine hormones and insulin sensitivity in human obesity. Diabetes Care 35:1105–1111.

- 15. van de Giessen E, de Bruin K, la Fleur SE, van den Brink W, Booij J (2012): Triple monoamine inhibitor tesofensine decreases food intake, body weight, and striatal dopamine D2/D3 receptor availability in dietinduced obese rats. Eur Neuropsychopharmacol 22:290–299.
- Guo J, Simmons WK, Herscovitch P, Martin A, Hall KD (2014): Striatal dopamine D2-like receptor correlation patterns with human obesity and opportunistic eating behavior. Mol Psychiatry 19:1078–1084.
- Karlsson HK, Tuominen L, Tuulari JJ, Hirvonen J, Parkkola R, Helin S, et al. (2015): Obesity is associated with decreased μ-opioid but unaltered dopamine D2 receptor availability in the brain. J Neurosci 35:3959–3965.
- Carpenter CL, Wong AM, Li Z, Noble EP, Heber D (2013): Association of dopamine D2 receptor and leptin receptor genes with clinically severe obesity. Obesity (Silver Spring) 21:E467–E473.
- Chen AL, Blum K, Chen TJH, Giordano J, Downs BW, Han D, et al. (2012): Correlation of the Taq1 dopamine D2 receptor gene and percent body fat in obese and screened control subjects: A preliminary report. Food Funct 3:40–48.
- 20. Barnard ND, Noble EP, Ritchie T, Cohen J, Jenkins DJA, Turner-McGrievy G, *et al.* (2009): D2 dopamine receptor Taq1A polymorphism, body weight, and dietary intake in type 2 diabetes. Nutrition 25:58–65.
- Comings DE, Gade R, MacMurray JP, Muhleman D, Peters WR (1996): Genetic variants of the human obesity (OB) gene: Association with body mass index in young women, psychiatric symptoms, and interaction with the dopamine D2 receptor (DRD2) gene. Mol Psychiatry 1:325–335.
- Noble EP, Noble RE, Ritchie T, Syndulko K, Bohlman MC, Noble LA, et al. (1994): D2 dopamine receptor gene and obesity. Int J Eat Disord 15:205–217.
- Hardman CA, Rogers PJ, Timpson NJ, Munafò MR (2014): Lack of association between DRD2 and OPRM1 genotypes and adiposity. Int J Obes (Lond) 38:730–736.
- 24. Beeler JA, Frazier CRM, Zhuang X (2012): Putting desire on a budget: Dopamine and energy expenditure, reconciling reward and resources. Front Integr Neurosci 6:49.
- 25. Di Chiara G, Tanda G, Cadoni C, Acquas E, Bassareo V, Carboni E (1998): Homologies and differences in the action of drugs of abuse and a conventional reinforcer (food) on dopamine transmission: An interpretative framework of the mechanism of drug dependence. Adv Pharmacol 42:983–987.
- Davis JF, Tracy AL, Schurdak JD, Tschöp MH, Lipton JW, Clegg DJ, Benoit SC (2008): Exposure to elevated levels of dietary fat attenuates psychostimulant reward and mesolimbic dopamine turnover in the rat. Behav Neurosci 122:1257–1263.
- Li Y, South T, Han M, Chen J, Wang R, Huang X-F (2009): High-fat diet decreases tyrosine hydroxylase mRNA expression irrespective of obesity susceptibility in mice. Brain Res 1268:181–189.
- Vucetic Z, Reyes TM (2010): Central dopaminergic circuitry controlling food intake and reward: Implications for the regulation of obesity. Wiley Interdiscip Rev Syst Biol Med 2:577–593.
- Chaput J-P, Klingenberg L, Rosenkilde M, Gilbert J-A, Tremblay A, Sjödin A (2011): Physical activity plays an important role in body weight regulation. J Obes2011.
- Elder SJ, Roberts SB (2007): The effects of exercise on food intake and body fatness: A summary of published studies. Nutr Rev 65:1–19.
- Chakravarthy MV, Booth FW (2004): Eating, exercise, and "thrifty" genotypes: Connecting the dots toward an evolutionary understanding of modern chronic diseases. J Appl Physiol 96:3–10.
- 32. Booth FW, Lees SJ (2007): Fundamental questions about genes, inactivity, and chronic diseases. Physiol Genomics 28:146–157.
- **33.** Hill JO, Wyatt HR, Reed GW, Peters JC (2003): Obesity and the environment: Where do we go from here? Science 299:853–855.
- 34. Levin BE, Dunn-Meynell AA (2004): Chronic exercise lowers the defended body weight gain and adiposity in diet-induced obese rats. Am J Physiol Regul Integr Comp Physiol 286:R771–R778.
- Warburton DER, Nicol CW, Bredin SSD (2006): Health benefits of physical activity: The evidence. CMAJ 174:801–809.

- Hawley JA, Holloszy JO (2009): Exercise:. It's the real thing! Nutr Rev 67:172–178.
- Powell KE, Blair SN (1994): The public health burdens of sedentary living habits: Theoretical but realistic estimates. Med Sci Sports Exerc 26:851–856.
- Berridge KC, Ho C-Y, Richard JM, DiFeliceantonio AG (2010): The tempted brain eats: Pleasure and desire circuits in obesity and eating disorders. Brain Res 1350:43–64.
- Davis C, Strachan S, Berkson M (2004): Sensitivity to reward: Implications for overeating and overweight. Appetite 42:131–138.
- Narayanaswami V, Thompson AC, Cassis LA, Bardo MT, Dwoskin LP (2013): Diet-induced obesity: Dopamine transporter function, impulsivity and motivation. Int J Obes (Lond) 37:1095–1103.
- Kok P, Roelfsema F, Frölich M, van Pelt J, Meinders AE, Pijl H (2006): Activation of dopamine D2 receptors lowers circadian leptin concentrations in obese women. J Clin Endocrinol Metab 91:3236–3240.
- Kim KS, Yoon YR, Lee HJ, Yoon S, Kim S-Y, Shin SW, et al. (2010): Enhanced hypothalamic leptin signaling in mice lacking dopamine D2 receptors. J Biol Chem 285:8905–8917.
- Pfaffly J, Michaelides M, Wang G-J, Pessin JE, Volkow ND, Thanos PK (2010): Leptin increases striatal dopamine D2 receptor binding in leptin-deficient obese (ob/ob) mice. Synapse 64:503–510.
- Baldo B, Sadeghian K, Basso A (2002): Effects of selective dopamine D1 or D2 receptor blockade within nucleus accumbens subregions on ingestive behavior and associated motor activity. Behav Brain Res 137:165–177.
- Salamone JD, Steinpreis RE, McCullough LD, Smith P, Grebel D, Mahan K (1991): Haloperidol and nucleus accumbens dopamine depletion suppress lever pressing for food but increase free food consumption in a novel food choice procedure. Psychopharmacology (Berl) 104:515–521.
- Salamone JD, Correa M, Mingote SM, Weber SM (2005): Beyond the reward hypothesis: Alternative functions of nucleus accumbens dopamine. Curr Opin Pharmacol 5:34–41.
- García-Tornadú I, Rubinstein M, Gaylinn BD, Hill D, Arany E, Low MJ, et al. (2006): GH in the dwarf dopaminergic D2 receptor knockout mouse: Somatotrope population, GH release, and responsiveness to GH-releasing factors and somatostatin. J Endocrinol 190:611–619.
- García-Tornadú I, Ornstein AM, Chamson-Reig A, Wheeler MB, Hill DJ, Arany E, et al. (2010): Disruption of the dopamine d2 receptor impairs insulin secretion and causes glucose intolerance. Endocrinology 151:1441–1450.
- Zachwieja JJ, Hendry SL, Smith SR, Harris RB (1997): Voluntary wheel running decreases adipose tissue mass and expression of leptin mRNA in Osborne-Mendel rats. Diabetes 46:1159–1166.
- Rhodes JS, Gammie SC, Garland T (2005): Neurobiology of mice selected for high voluntary wheel-running activity. Integr Comp Biol 45:438–455.
- Moran TH, Bi S (2006): Hyperphagia and obesity in OLETF rats lacking CCK-1 receptors. Philos Trans R Soc Lond B Biol Sci 361:1211–1218.
- Patterson CM, Levin BE (2008): Role of exercise in the central regulation of energy homeostasis and in the prevention of obesity. Neuroendocrinology 87:65–70.
- Patterson CM, Bouret SG, Dunn-Meynell AA, Levin BE (2009): Three weeks of postweaning exercise in DIO rats produces prolonged increases in central leptin sensitivity and signaling. Am J Physiol Regul Integr Comp Physiol 296:R537–R548.
- Meek TH, Eisenmann JC, Garland T (2010): Western diet increases wheel running in mice selectively bred for high voluntary wheel running. Int J Obes (Lond) 34:960–969.
- 55. Novak CM, Escande C, Burghardt PR, Zhang M, Barbosa MT, Chini EN, et al. (2010): Spontaneous activity, economy of activity, and resistance to diet-induced obesity in rats bred for high intrinsic aerobic capacity. Horm Behav 58:355–367.
- Klinker F, Hasan K, Paulus W, Nitsche MA, Liebetanz D (2013): Pharmacological blockade and genetic absence of the dopamine D2 receptor specifically modulate voluntary locomotor activity in mice. Behav Brain Res 242:117–124.

- Stice E, Yokum S, Zald D, Dagher A (2011): Dopamine-based reward circuitry responsivity, genetics, and overeating. Curr Top Behav Neurosci 6:81–93.
- Martínez-González MA, Martínez JA, Hu FB, Gibney MJ, Kearney J (1999): Physical inactivity, sedentary lifestyle and obesity in the European Union. Int J Obes Relat Metab Disord 23:1192–1201.
- Martínez JA, Kearney JM, Kafatos A, Paquet S, Martínez-González MA (1999): Variables independently associated with selfreported obesity in the European Union. Public Health Nutr 2: 125–133.
- Ladabaum U, Mannalithara A, Myer PA, Singh G (2014): Obesity, abdominal obesity, physical activity, and caloric intake in US adults: 1988 to 2010. Am J Med 127:717–727.e12.
- Qi L, Hu FB, Hu G (2008): Genes, environment, and interactions in prevention of type 2 diabetes: A focus on physical activity and lifestyle changes. Curr Mol Med 8:519–532.
- 63. Hu FB (2011): Diet and exercise for new-onset type 2 diabetes? Lancet 378:101–102.

- Héroux M, Janssen I, Lam M, Lee D-C, Hebert JR, Sui X, Blair SN (2010): Dietary patterns and the risk of mortality: Impact of cardiorespiratory fitness. Int J Epidemiol 39:197–209.
- 65. Healy GN, Wijndaele K, Dunstan DW, Shaw JE, Salmon J, Zimmet PZ, Owen N (2008): Objectively measured sedentary time, physical activity, and metabolic risk: The Australian Diabetes, Obesity and Lifestyle Study (AusDiab). Diabetes Care 31:369–371.
- Healy GN, Matthews CE, Dunstan DW, Winkler EAH, Owen N (2011): Sedentary time and cardio-metabolic biomarkers in US adults: NHANES 2003-06. Eur Heart J 32:590–597.
- Noble EE, Billington CJ, Kotz CM, Wang C (2014): Oxytocin in the ventromedial hypothalamic nucleus reduces feeding and acutely increases energy expenditure. Am J Physiol Regul Integr Comp Physiol 307:R737–R745.
- Zink AN, Perez-Leighton CE, Kotz CM (2014): The orexin neuropeptide system: Physical activity and hypothalamic function throughout the aging process. Front Syst Neurosci 8:211.
- Kotz CM, Wang C, Teske JA, Thorpe AJ, Novak CM, Kiwaki K, Levine JA (2006): Orexin A mediation of time spent moving in rats: Neural mechanisms. Neuroscience 142:29–36.